

A Proposal of Establishing Tribes for the Family Drosophilidae with Key to Tribes and Genera (Diptera)

TOYOHICHI OKADA

Gotokuji 2-30-18, Setagaya, Tokyo 154, Japan

ABSTRACT—Subdivision into tribes of the family Drosophilidae is attempted. Establishment of two tribes, Steganini and Leucophengini, for the subfamily Steganinae and five tribes, Microdrosophilini, Hypsiletryrini, Colocasiomyini, Dettopsomyini and Drosophilini, for Drosophilinae is proposed.

INTRODUCTION

By the end of 1984, 62 genera of the family Drosophilidae were recorded by Wheeler [1], including one fossil and five ungrouped into subfamilies. Since then, one genus, *Thyreoccephala*, was added by myself [2], which was, however, synonymized with *Mulgravea* Bock by myself [3], and two fossil genera, *Miomyia* and *Protoclymomyza*, by Grimaldi [4]. The genus *Drosophilella* Duda was recently synonymized with *Colocasiomyia* de Meijere by myself [5]. The genera *Atele-drosophila* Hardy and *Nudidrosophila* Hardy were treated as distinct genera according to Wheeler [6], although they were synonymized with *Drosophila* s. str. by Kaneshiro [7] because their females are not distinguishable from the latter.

To recognize tribe, a subordinate category of family or subfamily, is a general rule especially in large families of insects. No explicit subdivision of the family Drosophilidae has, however, been attempted before the present proposal, although an implicit subdivision into "Drosophiloids" and "Scaptomyzoids (as Scaptoids)" was made for endemic Hawaiian Drosophilidae by Throckmorton [8]. The inference that drosophilid taxonomy is highly developed might be accepted mostly at the level below species group.

METHODS

The descriptions of the tribes with diagnoses of

involving genera are given by a form of key. For establishing the tribes and constructing the key, following fourteen diagnostic characters ($n=14$: A-N) each divided into two states (0: presumed plesiomorph; 1, presumed apomorph) are taken into consideration, supplemented by some additional special characters.

Eye bare (A=0) or piled (a=1).

Arista plumose (B=0) or bare or pubescent (b =1).

Ocellars inside (C=0) or outside (c=1) ocellar triangle.

Postverticals present (D=0) or absent (d=1).

Carina undeveloped (E=0) or developed (e =1).

Anterior reclinate orbital large (F=0) or fine (f =1).

Posterior reclinate orbital nearer to inner vertical than to proclinate (G=0) or nearer to proclinate than to inner vertical (g=1).

Prescutellars developed (H=0) or absent (h =1).

Anterior dorsocentral nearer to scutellum than to suture (I=0) or nearer to suture than to scutellum (i=1).

Acrostichal hairs in more than 8 rows (J=0) or 8 or less than 8 rows (j=1).

Lateral scutellars convergent or parallel (K=0) or divergent (k=1).

Second costal break shallow (L=0) or deep (l =1).

Second costal lappet small (M=0) or large (m =1).

Discal and second basal cells separated (N=0)

Accepted June 21, 1988

Received April 23, 1988

TABLE 1. n(characters)×t(taxa) matrix

Taxa Genus	Characters													
	A	B	C	D	E	F	G	H	I	J	K	L	M	N
Steganinae														
1. Steganini														
[†] Electrophorica	0	0	0	0	0	0	0	0	0	1	1	0	0	0
Soederbomia	NC	0	NC	NC	0	0	0	0	0	0	0	NC	0	0
Pyrgometopha	NC	0	NC	NC	0	NC	0	NC	NC	NC	NC	0	0	0
Eostegana	0	0	0	0	1	0	0	0	0	0	1	0	0	0
Stegana	0	0	0	0	1	0	0	0	0	0	1	0	0	0
Amiota	0	0	0	0	1	0	0	0	0	0	1	0	0	0
Crincosia	0	1	1	9	1	9	0	0	0	0	0	1	0	0
Apenthecia	0	1	0	0	0	0	1	10	0	0	0	1	0	0
Mayagueza	0	1	0	0	0	0	0	0	0	0	1	0	0	0
Cacoxenus	0	1	0	0	1	0	0	0	0	0	1	0	0	0
Gitona	0	1	0	0	1	0	0	0	0	1	1	0	0	0
2. Leucophengini														2
Aclatoxenus	0	1	NC	0	1	0	0	0	0	0	0	0	0	1
Luzonimyia	0	1	13	1	0	1	0	0	0	0	0	1	0	1
Leucophenga	0	0	0	0	1	0	0	0	0	0	1	0	0	1
Paraleucophenga	0	0	0	0	1	0	0	0	0	0	1	0	0	1
Pseudastata	0	1	1	0	0	12	0	0	0	0	0	0	0	1
Trachyleucophenga	NC	0	16	NC	0	0	0	0	0	NC	NC	NC	0	1
Pararhinoleucophenga	0	0	0	0	0	0	0	0	0	0	0	1	0	1
Rhinoleucophenga	0	0	0	0	0	0	0	0	1	0	0	1	0	1
Drosophilinae														
1. Microdrosophilini														
Microdrosophila	1	0	0	0	0	1	1	1	1	1	1	1	0	1
2. Hypselothyrini														19-20
Hypselothryea	1	0	0	1	1	25	1	1	1	1	1	0	0	1
Tambourella	0	0	0	1	0	1	1	1	1	1	1	0	0	1
Sphaerogastrella	0	0	0	1	0	1	1	1	1	1	1	1	0	1
Liodesophila	0	0	0	0	1	1	1	1	1	1	1	1	0	1
Lissoccephala	1	0	0	0	1	27	1	1	1	1	1	1	0	1
Mulgravea	1	0	0	0	0	1	1	1	1	1	1	1	0	1
Paraliodesophila	1	0	NC	NC	0	NC	1	NC	NC	1	0	0	0	1

TABLE I. (continued)

Taxa Genus	A	B	C	D	E	F	G	H	I	J	K	L	M	N
3. Colocasiomyini														
<i>Calodrosophila</i>	1	0	1	0	1	1	1	1	0	1	0	0	0	1
<i>Colocasiomyia</i>	1	1	1	0	1	1	1	1	1	1	0	0	0	1
<i>Nesiodrosophila</i>	1	0	1	0	0	0	1	1	1	1	1	0	0	1
<i>Jeannelopsis</i>	1	0	1	0	0	0	1	1	1	1	1	0	0	1
4. Dettopsomyini				21									21-2	
<i>Mycodrosophila</i>	0	0	0	0	0	1	1	1	0	1	1	1	1	1
<i>Styloptera</i>	1	0	1	0	0	0	1	1	1	1	1	0	1	1
<i>Dettopsomyia</i>	1	0	0	0	1	1	1	1	1	1	1	1	1	1
<i>Paramycodrosophila</i>	1	0	0	0	0	1	1	1	0	1	1	1	1	1
5. Drosophilini												22	22	
<i>Dichadochaeta</i>	NC	1	NC	0	NC	1	1	NC	1	1	NC	0	0	1
<i>Baeodrosophila</i>	1	1	1	0	0	1	1	0	0	1	1	0	0	1
<i>Sphyrnoceps</i>	NC	0	NC	NC	0	1	1	1	NC	1	NC	0	0	1
<i>Cladochaeta</i>	0	0	0	0	0	1	1	1	0	1	0	0	0	1
<i>Miomyia</i>	0	0	0	NC	0	1	1	1	0	NC	NC	0	0	1
<i>Diathoneura</i>	1	0	0	0	0	1	1	1	1	1	0	0	0	1
<i>Neotanyastrella</i>	1	0	0	1	0	1	1	1	0	1	1	0	0	1
<i>Atelodrosophila</i>	1	0	NC	0	1	1	1	1	0	1	1	0	0	1
<i>Nudidrosophila</i>	1	0	1	0	0	0	0	0	0	1	1	0	0	1
<i>Zygothrica</i>	1	0	0	0	1	1	1	1	0	1	0	0	0	1
<i>Collessia</i>	1	0	0	0	1	1	1	1	0	1	1	0	0	1
<i>Phorticella</i>	1	0	0	0	1	1	1	1	0	1	1	0	0	1
<i>Grimshawomyia</i>	1	0	0	0	0	0	1	1	1	1	1	0	0	1
<i>Celidosoma</i>	NC	0	0	0	1	0	1	1	0	1	NC	0	0	1
<i>Protochymomyza</i>	0	0	0	0	1	0	1	1	0	1	1	0	0	1
<i>Chymomyza</i>	0	0	0	0	1	0	1	1	0	1	0	0	0	1
<i>Titanochaeta</i>	1	0	0	0	1	0	1	1	1	1	1	0	0	1
<i>Balara</i>	0	0	0	0	0	0	1	0	0	0	0	0	0	1
<i>Samoiaia</i>	1	0	0	0	0	0	1	1	1	1	1	0	0	1
<i>Scaptomyza</i>	1	0	0	0	0	0	1	1	1	1	1	0	0	1
<i>Marquesia</i>	NC	0	NC	0	0	0	1	1	0	1	NC	0	0	1
<i>Neorhynoleucophenga</i>	0	0	NC	0	0	0	1	1	NC	1	NC	0	1	1

TABLE I. (continued)

Genus	Characters													
	A	B	C	D	E	F	G	H	I	J	K	L	M	N
Zaprionus	1	0	0	0	0	0	0	1	0	1	0	0	0	1
Drosophila	1	0	0	0	0	0	1	1	0	1	0	0	0	1
Ungrouped genera														
Apachrochaeta	0	0	NC	0	0	NC	1	NC	0	NC	NC	0	0	1
Laccodosiphila	1	0	NC	0	0	0	0	0	0	NC	NC	0	0	1
Pseudocacoxenus	NC	1	NC	NC	0	NC	NC	0	1	NC	NC	0	0	1
Zapirothrica	NC	1	0	0	NC	1	1	1	0	NC	NC	0	0	1

0, Presumed plesiomorph; 1, presumed apomorph; NC, no comparison; †, fossil genus; numerical figure intercepting a longitudinal line, order of key couplet.





FIG. 1. Diagrams to show the key to tribes and genera of the family Drosophilidae. Alphabetical signs: Presumed plesiomorphous (large letters) and apomorphic states (small letters) of diagnostic characters; open and solid circles: plesiomorphs and apomorphs, respectively, appearing only once in the key; open and solid squares: plesiomorphs and apomorphs, respectively, repeatedly appearing in the key; numerical figures at the branching points of diagrams: orders of key couplets.



or confluent ($n=1$).

From the $n \times t$ matrix (Table 1), a key is constructed by means of "even dichotomous diagram method" [9]. The result is expressed also by key diagrams (Fig. 1), which may enable one to find the relationships of the tribes more easily.

Samples of some genera unknown to me were borrowed from various investigators and institutions: Dr. Ben Brugge, University of Amsterdam (*Colocasiomyia*). Dr. Don Colles, C.S.I.R.O., Canberra (*Luzonimyia*, *Crincosia*, *Balara*), Dr. K. Y. Kaneshiro, University of Hawaii (*Grimshawomyia*, *Titanochaeta*, *Samoaia*), and Dr. D. A. Grimaldi, American Museum of Natural History, New York (*Cladochaeta*, *Diathoneura*).

RESULTS

Family Drosophilidae

1. Posterior reclinate orbital nearer to inner vertical than to proclinate (G); prescutellars developed (H); tibiae and tarsi of mid and hind legs with dense rows of short bristles.Subfamily Steganinae.2.
- Posterior reclinate orbital nearer to proclinate than to inner vertical (g, excl. *Zaprimonius* s. str.); prescutellars usually absent (h); tibiae and tarsi of mid and hind legs without dense rows of short bristles....Subfamily Drosophilinae.19.

Subfamily Steganinae

2. Discal and second basal cells separated (N)....Tribe 1. Steganini.3.
- Discal and second basal cells confluent (n)....Tribe 2. Leucophengini12.

Tribe 1. Steganini Okada, n.

Type genus: *Stegana* Meigen

3. Arista usually plumose (B)4.
- Arista usually bare or pubescent (b).8.
4. Carina developed (e).5.
- Carina undeveloped (E).6.
5. Dorsocentrals in 2 pairs; R_{2+3} much waved; frons caudally protruded.....
.....Genus *Pyrgometopa* Kertész
- Dorsocentrals in 1 pair; R_{2+3} not waved;

- frons caudally not protruded.....5'.
- 5'. Acrostichal hairs in 8 or less than 8 rows (j).
.....Genus *Soederbomia* Hendel
- Acrostichal hairs in more than 8 rows (J).
.....Genus *Electrophortica* Hendel
- 6. Ac-index less than 4.0; wing maculated.
.....Genus *Eostegana* Hendel
- Ac-index more than 4.0; wing not maculated7.
- 7. Mid tibia usually with stout bristles above.
.....Genus *Stegana* Meigen
- Mid tibia without stout bristles above.
.....Genus *Amiota* Loew
- 8. Carina undeveloped (E).9.
- Carina developed (e).11.
- 9. Ocellars outside ocellar triangle (c); postverticals absent (d).Genus *Crincosia* Bock
- Ocellars inside ocellar triangle (C); postverticals present (D).10.
- 10. Anterior reclinate orbital fine (f).
.....Genus *Apenthecia* Tsacas
- Anterior reclinate orbital large (F).
.....Genus *Mayagueza* Wheeler
- 11. Acrostichal hairs in 8 or less than 8 rows (j).
.....Genus *Gitona* Meigen
- Acrostichal hairs in more than 8 rows (J).
.....Genus *Cacoxenus* Loew

Tribe 2. *Leucophengini* Okada, n.

Type genus: *Leucophenga* Mik

12. Carina undeveloped (E).13.
- Carina developed (e).16.
13. Arista pubescent (b).14.
- Arista plumose (B).15.
14. Lateral scutellars divergent or parallel (K); ocellae absent.
.....Genus *Aclotoxenus* Frauenfeld
- Lateral scutellars convergent (k); ocellars present, though minute.
.....Genus *Luzonimyia* Malloch
15. Excessive small scutellars absent.
.....Genus *Leucophenga* Mik
- Excessive small scutellars usually present.
.....Genus *Paraleucophenga* Hendel
16. Arista pubescent (b).
.....Genus *Pseudiastattata* Coquillett
- Arista plumose (B).17.

17. Frons punctured.
 Genus *Trachyleucophenga* Hendel
 — Frons not punctured.18.
18. Vein M curved to R_{4+5} distally.
 Genus *Pararhinoleucophenga* Hendel
 — Veins R_{4+5} and M divergent distally.
 Genus *Rhinoleucophenga* Hendel
- Subfamily Drosophilinae
19. Periorbit broadened anteriorly; second costal break deep (l); frons not glossy; ocellars inside ocellar triangle (C)....Tribe 1. Microdrosophilini.23.
 — Periorbit not broadened anteriorly20.
20. Frons glossy; second costal break shallow (L); ocellars inside ocellar triangle (C)....Tribe 2. Hypselothyrini.24.
- Frons not glossy.21.
21. Ocellars outside ocellar triangle (c); second costal break shallow (L)....Tribe 3. Colocasiomyini.30.
 — Ocellars usually inside ocellar triangle (C).22.
22. Second costal break deep (l; excl. *Styloptera*); second costal lappet usually large (m)....Tribe 4. Dettopsomyini.33.
 — Second costal break shallow (L); second costal lappet Small (M)....Tribe 5. Drosophilini.36.
- Tribe 1. Microdrosophilini Okada, n.
 Type genus: *Microdrosophila* Malloch
23. Monotypic.
 Genus *Microdrosophila* Malloch
- Tribe 2. Hypselothyrini Okada, n.
 Type genus: *Hypselothyrea* de Meijere
24. Postverticals absent (d).25.
 — Postverticals present (D).27.
25. Carina undeveloped (E); eye piled (a).
 Genus *Hypselothyrea* de Meijere
 — Carina developed (e); eye bare (A).26.
26. Lateral scutellars convergent (K) or parallel.
 Genus *Tambourella* Wheeler
 — Lateral scutellars divergent (k).
 Genus *Sphaerogastrella* Duda
27. Carina developed (e).28.
 — Carina undeveloped (E).29.
28. Eye bare (A); fore femur with a row of small spicules inside.Genus *Liodrosophila* Duda
 — Eye piled (a); fore femur without spicules inside.Genus *Lisocephala* Malloch
29. Lateral scutellars divergent (k).
 Genus *Mulgravea* Bock
 — Lateral scutellars convergent (K).
 Genus *Paraliiodrosophila* Duda
- Tribe 3. Colocasiomyini Okada, n.
 Type genus: *Colocasiomyia* de Meijere
30. Carina developed (e); anterior reclinata orbital fine (f); lateral scutellars convergent or parallel (K).31.
 — Carina undeveloped (E); anterior reclinata orbital large (F); lateral scutellars divergent (k).32.
31. Arista plumose (B); anterior dorsocentrals nearer to scutellum than to suture (I).
 Genus *Calodrosophila* Wheeler & Takada
 — Arista usually pubescent (b); anterior dorsocentrals nearer to suture than to scutellum (i).Genus *Colocasiomyia* de Meijere
32. Wing not crispy; R_{2+3} apically gently curved to costa.
 Genus *Nesiodrosophila* Wheeler & Takada
 — Wing crispy; R_{2+3} apically strongly curved to costa.Genus *Jeannelopsis* Séguay
- Tribe 4. Dettopsomyini Okada, n.
 Type genus: *Dettopsomyia* Lamb
33. Eye bare (A); anterior dorsocentral minute or absent. Genus *Mycodrosophila* Oldenberg
 — Eye piled (a); anterior dorsocentral large. 34.
34. Ocellars outside ocellar triangle (c); anterior reclinata orbital behind proclinate.
 Genus *Styloptera* Duda
 — Ocellars inside ocellar triangle (C); anterior reclinata orbital slightly before proclinate.35.
35. Carina developed (e); anterior dorsocentrals nearer to suture than to scutellum (i).
 Genus *Dettopsomyia* Lamb

- Carina undeveloped (E); anterior dorsocentrals nearer to scutellum than to suture (I). Genus *Paramycodrosophila* Duda
- Tribe 5. *Drosophilini* Okada, n.
Type genus: *Drosophila* Fallén
36. Anterior recline orbital fine (f). 37.
- Anterior recline orbital large (F). 48.
37. Arista pubescent (b) or with a few branches. 38.
- Arista plumose (B). 39.
38. Acrostichal hairs in 2 rows (j).
..... Genus *Dicladochaeta* Malloch
- Acrostichal hairs in more than 8 rows (J).
... Genus *Baeodrosophila* Wheeler & Takada
39. Carina undeveloped (E). 40.
- Carina developed (e). 44.
40. Head much broader than thorax.
..... Genus *Sphyrnoceps* de Meijere
- Head not broader than thorax. 41.
41. Eye bare (A). 42.
- Eye piled (a). 43.
42. Arista without lower branches.
..... Genus *Cladochaeta* Coquillett
- Arista with 2 lower branches.
..... Genus *Miomyia* Grimaldi
43. Postverticals absent (d); body slender.
..... Genus *Neotanygastrella* Duda
- Postverticals present (D); body thick.
..... Genus *Diathoneura* Duda
44. Arista inserted near apex of 3rd antennal joint. 45.
- Arista inserted near base of 3rd antennal joint. 46.
45. Arista without upper branches; orbita present.
..... Genus *Ateledosophila* Hardy (♂)
- Arista with upper branches; orbita absent.
..... Genus *Nudidrosophila* Hardy (♂)
46. Lateral scutellars convergent or parallel (K); proboscis long.
..... Genus *Zygothrica* Wiedemann
- Lateral scutellars divergent (k); proboscis not very long. 47.
47. Mesoscutum without white longitudinal stripes; R₄₊₅ and M divergent distally.
..... Genus *Collessia* Bock
- Mesoscutum with white longitudinal stripes; R₄₊₅ and M parallel distally.
..... Genus *Phorticella* Duda
48. Second antennal joint protruded anteriorly below. 49.
- Second antennal joint not protruded anteriorly below. 50.
49. Carina developed (e); anterior dorsocentrals nearer to suture than to scutellum (i); wing markings distinct.
..... Genus *Grimshawomyia* Hardy
- Carina undeveloped (E); anterior dorsocentrals nearer to scutellum than to suture (I); wing dark brown, basally hyaline.
..... Genus *Celidosoma* Hardy
50. Carina undeveloped (E). 51.
- Carina developed (e). 53.
51. Eye entirely or nearly bare (A); anterior dorsocentrals nearer to scutellum than to suture (I); anterior recline orbital before proclinate; arista with branches. 52.
- Eye piled (a); anterior dorsocentrals nearer to suture than to scutellum (i); lateral scutellars divergent (k); anterior recline orbital behind proclinate; arista without lower branches.
..... Genus *Titanochaeta* Knab
52. Lateral scutellars convergent (K).
..... Genus *Chymomyza* Czerny
- Lateral scutellars divergent (k).
..... Genus *Protachymomyza* Grimaldi
53. Prescutellars present (H); dorsocentrals in 3 pairs.
..... Genus *Balara* Bock
- Prescutellars usually absent (h); dorsocentrals in 2, 3, or 4 pairs. 54.
54. Anterior dorsocentral nearer to suture than to scutellum (i). 55.
- Anterior dorsocentral nearer to scutellum than to suture (I). 56.
55. Dorsocentrals in 3 pairs, body stout.
..... Genus *Samoia* Malloch
- Dorsocentrals in 2 pairs; body slender.
..... Genus *Scaptomyza* Hardy
56. Dorsocentrals in 4 pairs; lower margin of gena densely haired.
..... Genus *Malquessia* Malloch
- Dorsocentrals usually in 2 pairs; lower margin of gena not densely haired. 57.

57. Eye nearly bare (A).....
 Genus *Neorhinoleucophenga* Duda
 — Eye piled (a).....58.
58. Mesoscutum with white longitudinal stripes; posterior recline orbital usually nearer to inner vertical than to proclinate (G).
 Genus *Zaprionus* Coquillett
 — Mesoscutum without white longitudinal stripes; posterior recline orbital nearer to proclinate than to inner vertical (g).
 Genus *Drosophila* Fallén

Ungrouped Genera

1. Arista plumose (B).....2.
 — Arista pubescnt (b).....3.
2. Eye bare (A). ... Genus *Apachrochaeta* Duda
 — Eye piled (a). Genus *Laccodrosophila* Duda
3. Prescutellars present though weak (H); anterior dorsocentral nearer to suture than to scutellum (i). Genus *Pseudocacoxenus* Duda
 — Prescutellars absent (h); anterior dorsocentral nearer to scutellum than to suture (I).Genus *Zapriothrica* Wheeler

ACKNOWLEDGMENTS

My sincere thanks are due for the loan of material and valuable information to Dr. Ian R. Bock, La Trobe University, Victoria, Dr. Ben Brugge, University of Amsterdam, Dr. Don H. Colles, C. S. I. R. O., Canberra, Dr. David A. Grimaldi, American Museum of Natural History, New York, Dr. D. Elmo Hardy, University of Hawaii, Honolulu, Dr. Kenneth I. Kaneshiro, University of Hawaii, Honolulu, Dr. Wayne N. Mathis, Smithsonian Institution, Washington, and Dr. Francisca C. do

Val, University of São Paulo.

I deeply thank Dr. Marshall R. Wheeler, University of Texas, Austin, for reading manuscript.

REFERENCES

- Wheeler, M. R. (1986) Additions to the catalog or the World's Drosophilidae. In "The Genetics and Biology of *Drosophila*" Ed. by M. Ashburner, H. L. Carson and J. N. Thompson Jr., Academic Press, London, pp. 395-409.
- Okada, T. (1985) The genus *Lisocephala* Malloch and an allied new genus of Southeast Asia and New Guinea (Diptera, Drosophilidae). Kontyū, Tokyo, **53**: 335-345.
- Okada, T. (1987) Note on the genus *Mulgravea* (Diptera, Drosophilidae). Kontyū, Tokyo, **55**: 187.
- Grimaldi, D. A. (1987) Amber fossil Drosophilidae (Diptera), with particular reference to the Hispaniolan taxa. Am. Mus. Novit., **2880**: 1-23.
- Okada, T. (1988) Taxonomic note on *Colocasiomyia cristata* de Meijere with generic synonymy (Diptera, Drosophilidae). Proc. Jap. Soc. syst. Zool., **37**: 34-39.
- Wheeler, M. R. (1981) The Drosophilidae, taxonomic overview. In "The Genetics and Biology of *Drosophila*" Ed. by M. Ashburner, H. L. Carson and J. N. Thompson Jr. Academic Press, London, pp. 1-57.
- Kaneshiro, K. Y. (1976) A revision of generic concepts in the biosystematics of Hawaiian Drosophilidae. Proc. Hawaii Ent. Soc., **22**: 255-278.
- Throckmorton, L. H. (1966) The relationships of the endemic Hawaiian Drosophilidae. Univ. Texas Publ., **6615**: 335-396.
- Okada, T. (1977) The automatic construction of keys. Jap. Soc. syst. Zool, Circular, **50**: 6-11 (in Japanese).